

TRUSTED SINCE 1926
BONIDE[®]

BONIDE ANT FLEA & TICK KILLER GRANULES EPA Reg. No. 4-425 EPA Est. No. 4-NY-1

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Harmful if inhaled. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Environmental Hazards

This product is extremely toxic to fish. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in treated areas. Do not contaminate water when cleaning equipment or when disposing of equipment washwaters.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information.

If in eyes:	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed:	Call poison control center or doctor immediately for advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled:	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Read entire label before use.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Avoid contamination of food and feed. Utensils such as teaspoons and measuring cups should not be used for food purposes after use with insecticides. Do not allow children and pets to enter treated area until dust has settled.

STORAGE & DISPOSAL

STORAGE: Store product in original container in a locked storage area. Store and transport in an upright position.

DISPOSAL: *If empty* - Do not reuse this container. Place in trash or offer for recycling if available. *If partly filled* - Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

LAWNS

Three Easy Steps to use this Product

- Step #1:** Determine size of the job, measure the area to be treated.
- Step #2:** Use spreader setting chart to determine appropriate setting for your spreader.
- Step #3:** Treat as directed in Rate chart. Thorough coverage is important. Water in granules where indicated.

Note: The spreader settings listed are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust spreader setting, if necessary, in order to assure correct rate of application.

Ant, Flea & Tick Killer Granules

SPREADER SETTINGS (per 1,000 sq. ft.)

Spreader Make and Model	Grass or Dichondra Lawns 2 Pounds	White Grubs, Billbugs 2 1/2 pounds	St. Augustine Lawns 3 pounds
CYCLONE			
Broadcast	3 1/2 - 4	4 1/2	5 1/2
GREENVIEW			
Drop Spred III	3 - 3 1/2	4	5
Jet Spred III	4 - 4 1/2	5	6
REPUBLIC			
Drop EZ530	8 - 9	10	11
Broadcast EZ540	2 - 2 1/2	3	4
SCOTTS			
AG3	3 - 3 1/2	4	5
SG3	4 - 4 1/2	5	6
EG3	26 1/2 - 27	27 1/2	28
PF4	49 1/4 - 5 1/2	6	6 1/4
SEARS			
Drop 1935	5 - 5 1/2	6	7
Broadcast 1922	11 - 11 1/2	12	13

Treatment Areas	Pests	Use Rate	How to Use
Overall Lawn application for Grass and Dichondra Lawns	Ants, Armyworms, Brown Dog Ticks, Chinch Bugs, Clover Mites, Crickets, Earwigs, Fleas, Lawn Moths (Sod Webworms), Leafhoppers, Millipedes, Sowbugs	For application with other types of equipment, apply at the rate of 2 lbs. Per 1000 sq. ft. (an area 20' by 50') or 3.2 oz Per 100 sq. ft.	Apply with a fertilizer spreader. See spreader setting chart above. Apply when grass is dry. Water lawn lightly after application. Repeat as necessary up to once every 4 weeks.
	Billbugs, Grubs, Japanese Beetle Grubs, Dung Beetles (Ateanus Spretulus)	Apply 2 1/2 lbs. Per 1000 sq. ft. of lawn.	Water grass thoroughly after application. For Billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. For Grub control, except Dung Beetle, treat any time between late July and early October. For Dung Beetle Grub control, apply in spring or early summer when damage is first noticed. Water grass thoroughly with 1/4 to 1/2 inch of water after application. Treat again in late summer or early fall if a second generation occurs.
Spot Treatment	Ants	Apply 1 1/2 teaspoons over and around each anthill.	Water lightly after application. Repeat every 4 weeks or as new hills appear.
	Fire Ants	Apply 1 cup over, and 2 ft. around, each fire ant mound.	Water thoroughly, immediately after application. A minimum of 1 gallon of water should be applied to each mound area. Apply gently to avoid disturbing ants. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness. Note: For best results, apply in early morning or late evening hours. Treat new mounds as they appear.
Band Treatment around House Foundation	Ants, Fleas, Brown Dog Ticks, Millipedes, Clover Mites, Roaches, Spiders, Silverfish, Sowbugs (Pillbugs), Crickets, Earwigs	3.2 oz. for each 100 sq. ft. area	These pests are commonly found around or near foundations of houses, as well as in lawns. Certain of these pests may enter houses. Apply a 5-foot band of granules next to the house foundation. Distribute granules uniformly. Water lightly after application. Repeat as necessary up to once every 4 weeks.

BACK PANEL

For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.

Notice: Buyer assumes all responsibility for safety and use not in accordance with directions.