

EIGHT INSECT CONTROL Yard and Garden Granules

- Use in and around flower beds, vegetable gardens or ornamental plantings
- Use as a perimeter treatment around buildings
- Controls: Ants, Chinch Bugs, Crickets, Cutworms, Earwigs, Fleas, Grasshoppers, June Beetles, Sow Bugs, Ticks, White Grubs, and others

EPA Est. No. 4-NY-1

EPA Reg. No. 4-142

Buyers Guarantee Limited to Label Claims.

BONIDE PRODUCTS, INC.
6301 Sutliff Road
Oriskany, NY 13424

BN



62880107

1 lb.

**PRECAUTIONARY STATEMENTS
HAZARDOUS TO HUMANS
AND DOMESTIC ANIMALS**

CAUTION: MAY BE HARMFUL IF SWALLOWED. Avoid breathing dust. Do not take internally. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling this product. Do not eat, drink or use tobacco while working with this product. Wash hands and face before eating, drinking or using tobacco. Keep out of reach of children and domestic animals.

USER SAFETY REQUIREMENTS

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.
- Change clothing as soon as possible after use.
- Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- Do not allow children or pets to contact treated area until dust has settled.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water by cleaning equipment or disposal of wastes.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

If swallowed	Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
---------------------	---

If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
-------------------	---

If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
-------------------------------	---

If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
-------------------	---

NOTE TO PHYSICIAN: Carbaryl is a carbamate insecticide, which is a cholinesterase inhibitor. Atropine is antidotal. Narcotics and other sedatives should not be used. Further, drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended. In case of emergency call 1-800-UCC-HELP.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with treated surfaces. Allow dust to settle in treated areas before reentering. Do not repeat application more than once a week.

Not for use on plants grown for sale or other commercial use, or for other commercial use or for research purposes.

BARRIER PEST CONTROL AROUND BUILDINGS, IN AND AROUND FLOWER BEDS AND ORNAMENTAL PLANTINGS.

PERIMETER TREATMENT: For control of ants, ticks, carpenter ants, centipedes, crickets, earwigs, firebrats and silverfish, fleas, millipedes, scorpions and spiders; apply granules uniformly in a 6 foot band around the building. Apply granules at the rate of 6.6 ounces per 120 square feet (6 x 20 foot band). Restrict applications to shrub beds, foundation plantings, and lawn or soil areas immediately adjacent to the building. Lightly water treated area soon after application.

Ants: Spot treatment - apply 1 tablespoon of granules in and on each ant hill or nest.

Digger Wasps: Same as for ants except apply at dusk when insects are less active. Dust in the evening when the insect is in its burrow and cover with a mound of earth.

USE IN VEGETABLE GARDENS AND STRAWBERRIES.

To control ants, centipedes, crickets, cutworms, earwigs, grasshoppers, millipedes, sowbugs (pillbugs), springtails and ticks. Apply as directed in the following use rate table. Water lightly after application to release the chemical from the granules. Apply anytime before, during or after the growing season except as directed otherwise in the Use Rate Table for the listed crops. Do not use on crops other than those listed.

Crop	Use Rate
Asparagus	Application to spears: Maximum single rate application rate of 1 lb. per 1000 square foot application with a maximum seasonal application rate of 3 lb. per 1000 square feet. Application to Ferns: (after harvest of spears): maximum single application rate of 1 lb. per 1000 square feet with a maximum seasonal application rate of 2lb. per 1000 square feet. Retreatment interval 7 days when applied to ferns after harvest. Retreatment interval 3 days when application are to spears. PHI= 1 day The total maximum seasonal application rate for spears and ferns is 4.6 lb per 1000 square feet. The total number of applications is 3 prior to harvest with 2 additional applications to ferns only.
Beans, fresh and dried (snap, navy and kidney); pea, fresh and dried; cowpea, lentil	Maximum single application rate of .75 lb per 1000 square feet application with a maximum seasonal application rate of 3 lb. per 1000 square feet. Retreatment interval = 7 days. PHI=3 days (succulents) PHI=21 days (dried) PHI=14 days (forage) PHI=21 days (hay)

Crop	Use Rate
Carrots, garden beets (roots), parsnip, rutabaga, radish, horseradish, salsify	Maximum single application rate of 1 lb. per 1000 square feet application with a maximum seasonal application rate of 3 lb. per 1000 square feet. Retreatment interval =7 days PHI=7 days Total number of applications/crop can be 6 (some of which must be at lower than maximum single application rate)
Sugar beets	Maximum single application rate of .75 lb. per 1000 square feet application with a maximum seasonal application rate of 1.5 lb. per 1000 square feet. Retreatment interval=14 days PHI=28 days (roots, forage)
Brussel sprouts, broccoli, cabbage, cauliflower, Chinese cabbage, collards, kale, kohlrabi	Maximum single application rate of 1 lb. per 1000 square feet application with a maximum seasonal application rate of 3 lb. per 1000 square feet. Retreatment interval=7 days PHI=3 days (brussel sprouts, broccoli, cabbage, kohlrabi, cauliflower) PHI=14 days (Chinese cabbage, kale, collards) Total number of applications/crop can be 4 (some of which must be at lower than maximum single application rate).
Corn (sweet)	Maximum single application rate of 1 lb. per 1000 square feet application with a maximum seasonal application rate of 8 lb. per 1000 square feet. Retreatment = 3 days PHI = 2 days (ears) PHI = 14 days (forage) PHI = 48 days (fodder)
Cucumber, melon, squash	Maximum single application rate of .5 lb. per 1000 square feet application with a maximum seasonal application rate of 3 lb. per 1000 square feet. Retreatment=7 days PHI=3 days
Lettuce, head and leaf, endive, celery, dandelion, Parsley, spinach, Swiss chard, garden beets (tops)	Maximum single application rate of 1 lb. per 1000 square feet application with a maximum seasonal application rate of 3 lb. per 1000 square feet. Retreatment=7 days PHI=14 to 15 days Total number of applications/crop can be 5 (some of which must be at a lower than maximum single application rate)

Crop	Use Rate
Potato	Maximum single application rate of 1 lb. per 1000 square feet application with a maximum seasonal application rate of 3 lb. per 1000 square feet. Retreatment = 7 days PHI = 7 days Total number of applications/crop can be 6 (some of which must be at a lower than maximum single application rate).
Strawberry	Maximum single application rate of 1 lb. per 1000 square feet application with a maximum seasonal application rate of 5 lb. per 1000 square feet. Retreatment = 7 days PHI = 7 days
Tomato, pepper, eggplant	Maximum single application rate of 1 lb. per 1000 square feet application with a maximum seasonal application rate of 4 lb. per 1000 square feet. Retreatment = 7 days PHI = 3 days Total number of applications/crop can be 7 (some of which must be at lower than maximum single application rate).

LAWNS, RECREATIONAL AND ORNAMENTAL TURF AREAS

For control of turfgrass pests, apply granules uniformly to the turf grass. For optimum control, follow specific directions for proper watering or irrigation after application. Do not allow public use of treated area during application. Before each application mow lawn, remove clipping and water thoroughly.

INSECTS	POUNDS PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
Ants, Armyworm, Centipedes, Chiggers, Chinch bugs, Cutworms, Earwigs, Essex skipper, European chafer, Fall armyworm, Fiery skipper, Grasshoppers, June beetles, Leafhoppers, Lucerne moth, Millipedes, Adult mosquitoes, Sowbugs, Spittlebugs, Springtails, Ticks, Yellowstriped armyworm, Lawn Moth (Sod webworm)	2.75 to 3.50 (20 lbs./5715 to 7270 sq. ft.)	For general insect control, apply granules uniformly when grass is dry. Water lawn lightly after application. Repeat as necessary to control new infestations.

IMPORTED FIRE ANT CONTROL - For use as a mound treatment to control imported fire ants, apply 1.5 to 3.0 oz. by weight over and around each fire ant mound. Irrigate mound thoroughly immediately after treatment to remove the chemical from the granules and flush it down into the mound. A minimum of 1 gallon of water should be applied to each mound, but do not use more than 2 gallons of water as flooding of the mound may result. Apply water gently to avoid disturbing ants and causing migration before chemical can be effective. For best results, apply in the early morning or late evening hours when ants are least active. Treat new mounds as they appear.

STORAGE AND DISPOSAL

Storage: Store unused product in original container only in cool, dry area out of reach of children and animals. Do not store in areas where temperatures frequently exceed 100°F.

Disposal: *If empty* - Do not reuse this container. Place in trash or offer for recycling if available. If partly filled - Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE OF DISCLAIMER: BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL INJURY, LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS, OR CAUTIONS.